Award Number: DAMD17-02-1-0683

TITLE: Breast Cancer Prevention by Inducing Apoptosis in DCIS

Using Breast Ductal Lavage

PRINCIPAL INVESTIGATOR: Patrick P. Koty, Ph.D.

CONTRACTING ORGANIZATION: University of Pittsburgh

Pittsburgh, PA 15260

REPORT DATE: September 2003

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

Form Approved

REPORT DOCUMENTATION PAGE

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY	2. REPORT DATE	3. REPORT TYPE AND DATES COVERED			
(Leave blank)	September 2003	Annual(1 Sep 2002 - 31 Aug 2003)			
4. TITLE AND SUBTITLE		——————————————————————————————————————	5. FUNDING N	IUMBERS	
Breast Cancer Prevention by Inducing Apoptosis in Do		is in DCIS	DAMD17-02	-1-0683	
Using Breast Ductal Lava	ıge				
6. AUTHOR(S)			-		
Patrick P. Koty, Ph.D.					
	***************************************				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER		
University of Pittsburgh Pittsburgh, PA 15260			REPORT NO	INBER	
ricessargii, FA 15200					
E-Mail: koty@pitt.edu					
9. SPONSORING / MONITORING			10. SPONSORING / MONITORING		
AGENCY NAME(S) AND ADDRESS	(ES)			REPORT NUMBER	
U.S. Army Medical Resear		and			
Fort Detrick, Maryland	21702-5012				
			•		
11. SUPPLEMENTARY NOTES			<u> </u>		
12a. DISTRIBUTION / AVAILABILITY S				12b. DISTRIBUTION CODE	
Approved for Public Rele	ease; Distribution Un	limited			
13. ABSTRACT (Maximum 200 Words	s)			1	
				•	
				second highest cause of cancer-	
related deaths for women in the Ur					
cancer incidence but increases the					
lesions that are estrogen receptor (F					
receptor. We hypothesize that pro ER-positive and ER-negative cells	grammed cell death is dysrego	ulated in premailignant	and malignant t	oreast cells which permits both	
molecular genetic-based agent may	to avoid cen deam. We mu	end to investigate whe	tment to induce	cell death in both FR-positive	
and ER-negative cells. Ultimately					

14. SUBJECT TERMS Chemoprevention, apopt in situ, bcl-2 gene fa	15. NUMBER OF PAGES 6		
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	Unlimited

ductal lobe of these high risk individuals thus eliminating any potential for tamoxifen-induced uterine cancers.

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. 239-18 298-102

# **Table of Contents**

Cover	
SF 298	1
Table of Contents	2
Introduction	3
Body	3
Key Research Accomplishments	5
Reportable Outcomes	5
Conclusions	5
References	5
Appendices	5

### TITLE:

Breast Cancer Prevention by Inducing Apoptosis in DCIS Using Breast Ductal Lavage

# Introduction:

Although great strides have been made in breast cancer screening and treatment, it remains the second highest cause of cancer-related deaths for women in the United States. Current prevention has focused on oral administration of tamoxifen which appears to decrease breast cancer incidence but increases the risk for secondary uterine cancer. In addition, tamoxifen may not be effective in preventing those lesions that are estrogen receptor (ER) negative based on its primary function of suppressing cell proliferation by blocking the estrogen receptor. We hypothesize that programmed cell death (PCD) is dysregulated in premalignant and malignant breast cells which permits both ER-positive and ER-negative cells to avoid cell death. We intend to investigate whether treating premalignant breast cells with a molecular genetic-based agent (antisense bcl-2/bcl-xL oligonucleotide) may be effective alone or in concert with tamoxifen treatment to induce cell death in both ER-positive and ER-negative cells. Ultimately, we envision using the newly developed technique of breast ductal lavage to not only screen women for increased risk in developing breast cancer, which is currently being performed, but to also use this technique to delivery genetic-based preventive agents and/or tamoxifen directly to the breast ductal lobe of these high risk individuals thus eliminating any potential for tamoxifen-induced uterine cancers.

# BODY:

The following is the approved Statement of Work for the project entitled "Breast Cancer Prevention by Inducing Apoptosis in DCIS Using Breast Ductal Lavage." When the Award was granted, the terms indicated that no patient samples could be obtained until the DoD's Human Subject Protection Office approved this project for the Use of Human Anatomical Substances. This still has not occurred, even though, this project received IRB approval on October 14, 2002 from the institution (Magee Womens Hospital, Pittsburgh PA) for which the samples are to be obtained. The documentation requested by the Human Subject Protection Office was sent to Ms. Catherine Smith on October 21, 2002 then transferred to Ms. Robin Dillner on November 5, 2002 but no decision on approval status was ever given. On April 1, 2003 I moved my researched laboratory from the University of Pittsburgh, Pittsburgh PA (who only released the funds for this project on August 13, 2003) to Wake Forest University School of Medicine, Winston-Salem NC. Since being here, I submitted (August 18, 2003) a modification of my IRB approved protocol at Magee Womens Hospital, Pittsburgh PA to send any future patient samples to my research laboratory at Wake Forest University School of Medicine. It has been approved but some clarifications need to be addressed by October 13, 2003. Although I have submitted an IRB application for review by the IRB at Wake Forest University School of Medicine, they are waiting for Magee Womens Hospital's IRB approval prior to approving at Wake Forest. Ms. Dillner indicated on September 16, 2003 that she would not be able to review the protocol that was initially sent and I would need to resubmit an application to the DoD's Human Subject Protection Office which

needs to include Wake Forest University School of Medicine IRB approval letter. Therefore, I remain without DoD's Human Subject Protection Office's approval for this project for the Use of Human Anatomical Substances and can not perform any research until I receive such approval as indicated on the terms of the subject award.

# Statement of Work

- Task 1. Determine expression pattern of the PCD regulatory genes bcl-2, bax, and bcl-xL in primary DCIS cultures (Months 1-6):
  - a. Protein analysis of Bcl-2, Bax, and Bcl-xL using Western blotting and immunohistochemical staining.
- Task 2. Determine whether down-regulation by genetic manipulation of the antiapoptotic genes bcl-2 and/or bcl-xL alone or in conjunction with physiological preventive doses of tamoxifen has the highest induction of PCD in primary DCIS cell cultures (Months 6-18):
  - a. Treatment with antisense and control oligonucleotides and/or tamoxifen.
  - b. Protein analysis of Bcl-2, Bax, and Bcl-xL using Western blotting and immunohistochemical staining.
  - c. Quantify mRNA for bcl-2 or bcl-xL using a PCR-based assay.
  - d. Determine effect of treatment on programmed cell death markers using assays for DNA fragmentation and caspase activation.
- Task 3. Determine expression pattern of the PCD regulatory genes bcl-2, bax, and bcl-xL in cells obtained by breast ductal lavage (Months 18-24):
  - a. Protein analysis of Bcl-2, Bax, and Bcl-xL using Western blotting and immunohistochemical staining.
- Task 4. Determine whether down-regulation by genetic manipulation of the anti-apoptotic genes bcl-2 and/or bcl-xL alone or in conjunction with physiological preventive doses of tamoxifen has the highest induction of PCD in cells from breast ductal lavages (Months 24-36).
  - a. Treatment with antisense and control oligonucleotides and/or tamoxifen.
  - b. Protein analysis of Bcl-2, Bax, and Bcl-xL using Western blotting and immunohistochemical staining.
  - c. Quantify mRNA for bcl-2 or bcl-xL using a PCR-based assay.
  - d. Determine effect of treatment on programmed cell death markers using assays for DNA fragmentation and caspase activation.

PI: Koty, Patrick P.

DAMD17-02-1-0683

KEY RESEARCH ACCOMPLISHMENTS: N/A

REPORTABLE OUTCOMES: N/A

Conclusions: N/A

REFERENCES: N/A

APPENDICES: N/A